



Complete Summary

GUIDELINE TITLE

Medical management of ectopic pregnancy.

BIBLIOGRAPHIC SOURCE(S)

American College of Obstetricians and Gynecologists (ACOG). Medical management of ectopic pregnancy. Washington (DC): American College of Obstetricians and Gynecologists (ACOG); 2008 Jun. 7 p. (ACOG practice bulletin; no. 94). [29 references]

GUIDELINE STATUS

This is the current release of the guideline.

This guideline updates a previous version: American College of Obstetricians and Gynecologists (ACOG). Medical management of tubal pregnancy. Washington (DC): American College of Obstetricians and Gynecologists (ACOG); 1998 Dec. 7 p. (ACOG practice bulletin; no. 3).

COMPLETE SUMMARY CONTENT

SCOPE
METHODOLOGY - including Rating Scheme and Cost Analysis
RECOMMENDATIONS
EVIDENCE SUPPORTING THE RECOMMENDATIONS
BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS
CONTRAINDICATIONS
QUALIFYING STATEMENTS
IMPLEMENTATION OF THE GUIDELINE
INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT
CATEGORIES
IDENTIFYING INFORMATION AND AVAILABILITY
DISCLAIMER

SCOPE

DISEASE/CONDITION(S)

Tubal ectopic pregnancy

GUIDELINE CATEGORY

Counseling
Diagnosis

Management
Treatment

CLINICAL SPECIALTY

Obstetrics and Gynecology

INTENDED USERS

Physicians

GUIDELINE OBJECTIVE(S)

- To aid practitioners in making decisions about appropriate obstetric and gynecologic care
- To review the risks and benefits of the use of methotrexate in the management of ectopic pregnancy

TARGET POPULATION

Pregnant women presenting with signs and symptoms of ectopic pregnancy

INTERVENTIONS AND PRACTICES CONSIDERED

Diagnosis

1. Clinical signs, physical symptoms
2. Serial human chorionic gonadotropin (hCG) levels
3. Transvaginal ultrasonography
4. Serum progesterone levels

Management/Treatment

1. Determining if a patient is a candidate for methotrexate therapy
2. Methotrexate
 - Single-dose regimen protocol
 - Two-dose regimen protocol
 - Fixed multidose regimen protocol
3. Surveillance after methotrexate therapy
 - Serial hCG levels
4. Patient counseling and education concerning side effects of methotrexate
5. Expectant management of ectopic pregnancy

MAJOR OUTCOMES CONSIDERED

- Sensitivity and specificity of diagnostic tests
- Incidence of heterotopic pregnancy
- Success rate of methotrexate regimen
- Methotrexate-associated morbidity
- Tubal preservation
- Tubal patency

- Repeat ectopic pregnancy
- Future pregnancies
- Patient satisfaction

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Hand-searches of Published Literature (Primary Sources)
 Hand-searches of Published Literature (Secondary Sources)
 Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

The MEDLINE database, the Cochrane Library, and the American College of Obstetricians and Gynecologists' (ACOG's) own internal resources and documents were used to conduct a literature search to locate relevant articles published between January 1985 and May 2007. The search was restricted to articles published in the English language. Priority was given to the articles reporting results of original research although review articles and commentaries also were consulted. Abstracts of research presented at symposiums and scientific conferences were not considered adequate for inclusion in this document. Guidelines published by organizations or institutions such as the National Institutes of Health and the American College of Obstetricians and Gynecologists were reviewed, and additional studies were located by reviewing bibliographies of identified articles.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Weighting According to a Rating Scheme (Scheme Given)

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Studies were reviewed and evaluated for quality according to the method outlined by the U.S. Preventive Services Task Force.

I Evidence obtained from at least one properly designed randomized controlled trial.

II-1 Evidence obtained from well-designed controlled trials without randomization.

II-2 Evidence obtained from well-designed cohort or case-control analytic studies, preferably from more than one center or research group.

II-3 Evidence obtained from multiple time series with or without the intervention. Dramatic results in uncontrolled experiments also could be regarded as this type of evidence.

III Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.

METHODS USED TO ANALYZE THE EVIDENCE

Review of Published Meta-Analyses
Systematic Review

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Expert Consensus

DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS

Analysis of available evidence was given priority in formulating recommendations. When reliable research was not available, expert opinions from obstetrician-gynecologists were used. See also the "Rating Scheme for the Strength of Recommendations" field regarding Grade C recommendations.

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Based on the highest level of evidence found in the data, recommendations are provided and graded according to the following categories:

Level A — Recommendations are based on good and consistent scientific evidence.

Level B — Recommendations are based on limited or inconsistent scientific evidence.

Level C — Recommendations are based primarily on consensus and expert opinion.

COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

METHOD OF GUIDELINE VALIDATION

Internal Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Practice Bulletins are validated by two internal clinical review panels composed of practicing obstetrician-gynecologists generalists and sub-specialists. The final guidelines are also reviewed and approved by the American College of Obstetricians and Gynecologists (ACOG) Executive Board.

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

The grades of evidence (I-III) and levels of recommendations (A-C) are defined at the end of "Major Recommendations" field.

The following conclusion is based on good and consistent evidence (Level A):

- In comparing systemic methotrexate with tube-sparing laparoscopic surgery, randomized trials have shown no difference in overall tubal preservation, tubal patency, repeat ectopic pregnancy, or future pregnancies.

The following recommendations and conclusions are based on limited or inconsistent evidence (Level B):

- An increase in serum human chorionic gonadotropin (hCG) of less than 53% in 48 hours confirms an abnormal pregnancy.
- With an hCG level of 5,000 mIU/mL or higher, multiple doses of methotrexate may be appropriate.
- Methotrexate can be considered in those women with a confirmed, or high clinical suspicion of, ectopic pregnancy who are hemodynamically stable with an unruptured mass.
- Failure of the hCG level to decrease by at least 15% from day 4 to day 7 after methotrexate administration is considered treatment failure requiring therapy with either additional methotrexate administration or surgical intervention.
- Post-treatment hCG levels should be monitored until a nonpregnancy level is reached.

The following conclusion is based primarily on consensus and expert opinion (Level C):

- If the initial hCG level is less than 200 mU/mL, 88% of patients experience spontaneous resolution.

Definitions:

Grades of Evidence

I Evidence obtained from at least one properly designed randomized controlled trial.

II-1 Evidence obtained from well-designed controlled trials without randomization.

II-2 Evidence obtained from well-designed cohort or case-control analytic studies, preferably from more than one center or research group.

II-3 Evidence obtained from multiple time series with or without the intervention. Dramatic results in uncontrolled experiments could also be regarded as this type of evidence.

III Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.

Levels of Recommendations

Level A — Recommendations are based on good and consistent scientific evidence.

Level B — Recommendations are based on limited or inconsistent scientific evidence.

Level C — Recommendations are based primarily on consensus and expert opinion.

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

The type of supporting evidence is identified and graded for each recommendation (see "Major Recommendations").

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

Improved treatment of tubal ectopic pregnancies with methotrexate

POTENTIAL HARMS

Adverse Effects Associated with Methotrexate Treatment

- Because methotrexate affects all rapidly dividing tissues within the body, including bone marrow, the gastrointestinal mucosa, and the respiratory epithelium, it should not be given to women with blood dyscrasias or active gastrointestinal and respiratory disease. Methotrexate is directly toxic to the

hepatocytes and is cleared from the body by renal excretion; therefore, it should not be used in women with liver or kidney disease.

- Methotrexate morbidity usually is dose and treatment duration dependent. Because methotrexate affects rapidly dividing tissues, gastrointestinal side effects, such as nausea, vomiting, and stomatitis, are the most common. Therefore, women treated with methotrexate should be advised not to use alcohol and nonsteroidal anti-inflammatory drugs (NSAIDs). Elevation of liver enzymes usually is seen only with multidose regimens and resolves after discontinuing methotrexate use or increasing the rescue dose of folinic acid. Alopecia is a rare side effect with the doses used to treat ectopic pregnancy.
- It is not unusual for women treated with methotrexate to experience abdominal pain 2–3 days after administration, presumably from the cytotoxic effect of the drug on the trophoblast tissue, causing tubal abortion.

CONTRAINDICATIONS

CONTRAINDICATIONS

Absolute Contraindications to Methotrexate Therapy

- Breastfeeding
- Overt or laboratory evidence of immunodeficiency
- Alcoholism, alcoholic liver disease, or other chronic liver disease
- Preexisting blood dyscrasias, such as bone marrow hypoplasia, leukopenia, thrombocytopenia, or significant anemia
- Known sensitivity to methotrexate
- Active pulmonary disease
- Peptic ulcer disease
- Hepatic, renal, or hematologic dysfunction

Relative Contraindications to Methotrexate Therapy

- Gestational sac larger than 3.5 cm
- Embryonic cardiac motion

QUALIFYING STATEMENTS

QUALIFYING STATEMENTS

These guidelines should not be construed as dictating an exclusive course of treatment or procedure. Variations in practice may be warranted based on the needs of the individual patient, resources, and limitations unique to the institution or type of practice.

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

An implementation strategy was not provided.

IMPLEMENTATION TOOLS

Audit Criteria/Indicators
Foreign Language Translations
Patient Resources

For information about [availability](#), see the "Availability of Companion Documents" and "Patient Resources" fields below.

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Getting Better

IOM DOMAIN

Effectiveness
Patient-centeredness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

American College of Obstetricians and Gynecologists (ACOG). Medical management of ectopic pregnancy. Washington (DC): American College of Obstetricians and Gynecologists (ACOG); 2008 Jun. 7 p. (ACOG practice bulletin; no. 94). [29 references]

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

1998 Dec (revised 2008 Jun)

GUIDELINE DEVELOPER(S)

American College of Obstetricians and Gynecologists - Medical Specialty Society

SOURCE(S) OF FUNDING

American College of Obstetricians and Gynecologists (ACOG)

GUIDELINE COMMITTEE

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Not stated

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

Not stated

GUIDELINE STATUS

This is the current release of the guideline.

This guideline updates a previous version: American College of Obstetricians and Gynecologists (ACOG). Medical management of tubal pregnancy. Washington (DC): American College of Obstetricians and Gynecologists (ACOG); 1998 Dec. 7 p. (ACOG practice bulletin; no. 3).

GUIDELINE AVAILABILITY

Electronic copies: None available

Print copies: Available for purchase from the American College of Obstetricians and Gynecologists (ACOG) Distribution Center, PO Box 4500, Kearneysville, WV 25430-4500; telephone, 800-762-2264, ext. 192; e-mail: sales@acog.org. The ACOG Bookstore is available online at the [ACOG Web site](#).

AVAILABILITY OF COMPANION DOCUMENTS

Proposed performance measures are included in the original guideline document.

PATIENT RESOURCES

The following is available:

- Ectopic pregnancy. Atlanta (GA): American College of Obstetricians and Gynecologists (ACOG); 2002.

Electronic copies: Available from the [American College of Obstetricians and Gynecologists \(ACOG\) Web site](#). Copies are also available in Spanish.

Print copies: Available for purchase from the American College of Obstetricians and Gynecologists (ACOG) Distribution Center, PO Box 4500, Kearneysville, WV 25430-4500; telephone, 800-762-2264, ext. 192; e-mail: sales@acog.org. The ACOG Bookstore is available online at the [ACOG Web site](#).

Please note: This patient information is intended to provide health professionals with information to share with their patients to help them better understand their health and their diagnosed disorders. By providing access to this patient information, it is not the intention of NGC to provide specific medical

advice for particular patients. Rather we urge patients and their representatives to review this material and then to consult with a licensed health professional for evaluation of treatment options suitable for them as well as for diagnosis and answers to their personal medical questions. This patient information has been derived and prepared from a guideline for health care professionals included on NGC by the authors or publishers of that original guideline. The patient information is not reviewed by NGC to establish whether or not it accurately reflects the original guideline's content.

NGC STATUS

This NGC summary was completed by ECRI on January 14, 2005. This summary was updated by ECRI Institute on July 29, 2008. The updated information was verified by the guideline developer on August 20, 2008.

COPYRIGHT STATEMENT

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

Date Modified: 11/3/2008

